

Quarks and gravitation

G. Quznetsov
gunn@chelcom.ru, gunn@mail.ru

February 10, 2008

Abstract

Turnings of the colors quarks mass space make curved the events time-space.

=====

In accordance with [1] the fermion moving equation has got the following form:

$$\begin{pmatrix} \beta^{[0]} (\partial_0 + i\Theta_0 + i\Upsilon_0\gamma^{[5]} - iM_0\gamma^{[0]} - iM_4\beta^{[4]}) \\ +\beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1\gamma^{[5]} + M_{\zeta,0}\beta^{[4]} - M_{\zeta,4}\gamma^{[0]}) \\ +\beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2\gamma^{[5]} + M_{\eta,0}\beta^{[4]} + M_{\eta,4}\gamma^{[0]}) \\ +\beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3\gamma^{[5]} - M_{\theta,0}\beta^{[4]} - M_{\theta,4}\gamma^{[0]}) \end{pmatrix} \varphi = 0. \quad (1)$$

Here [2]:

$$\beta^{[1]} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix}, \beta^{[2]} = \begin{bmatrix} 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \end{bmatrix},$$

$$\beta^{[3]} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

are the diagonal elements of the light Clifford pentad;

$$\gamma^{[0]} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, \beta^{[4]} = \begin{bmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{bmatrix}$$

are the antidiagonal (mass) elements of that pentad;

Θ_k and $\Upsilon_0\gamma^{[5]}$ with $k \in \{0, 1, 2, 3\}$ are fields, forming the electroweak gauge fields [4];

M_0, M_4 are lepton's mass numbers[5];

$M_{\zeta,0}, M_{\zeta,4}$ are red down and up quark's mass numbers;

$M_{\eta,0}, M_{\eta,4}$ are green down and up quark's mass numbers;

$M_{\theta,0}, M_{\theta,4}$ are blue down and up quark's mass numbers.

And

$$\gamma^{[5]} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}; \beta^{[0]} = - \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = -1_4 \quad [3]$$

The equation (1) can be reformulated as the following [1]:

$$\frac{1}{c} \partial_t \varphi + (i\Theta_0 + i\Upsilon_0 \gamma^{[5]}) \varphi = \begin{pmatrix} \beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ + \beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ + \beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} - \\ - iM_{\zeta,0} \gamma_{\zeta}^{[0]} + iM_{\zeta,4} \zeta^{[4]} - \\ - iM_{\eta,0} \gamma_{\eta}^{[0]} - iM_{\eta,4} \eta^{[4]} + \\ + iM_{\theta,0} \gamma_{\theta}^{[0]} + iM_{\theta,4} \theta^{[4]} \end{pmatrix} \varphi. \quad (2)$$

Here [2]:

$$\gamma_{\zeta}^{[0]} = \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix}, \zeta^{[4]} = \begin{bmatrix} 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ -i & 0 & 0 & 0 \end{bmatrix}$$

are the antidiagonal (mass) elements of red pentad;

$$\gamma_{\eta}^{[0]} = \begin{bmatrix} 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \end{bmatrix}, \eta^{[4]} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

are the antidiagonal (mass) elements of green pentad;

$$\gamma_{\theta}^{[0]} = \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}; \theta^{[4]} = \begin{bmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{bmatrix}$$

are the antidiagonal (mass) elements of blue pentad.

The quarks mass members of this equation form the following matrix sum:

$$\begin{aligned} \widehat{M} &\stackrel{def}{=} \begin{aligned} &-M_{\zeta,0} \gamma_{\zeta}^{[0]} + M_{\zeta,4} \zeta^{[4]} - \\ &-M_{\eta,0} \gamma_{\eta}^{[0]} - M_{\eta,4} \eta^{[4]} + \\ &+ M_{\theta,0} \gamma_{\theta}^{[0]} + M_{\theta,4} \theta^{[4]} \end{aligned} \\ &= \begin{bmatrix} 0 & 0 & -M_{\theta,0} & M_{\zeta,0} - iM_{\eta,0} \\ 0 & 0 & M_{\zeta,0} + iM_{\eta,0} & M_{\theta,0} \\ -M_{\theta,0} & M_{\zeta,0} - iM_{\eta,0} & 0 & 0 \\ M_{\zeta,0} + iM_{\eta,0} & M_{\theta,0} & 0 & 0 \end{bmatrix} \\ &+ i \begin{bmatrix} 0 & 0 & -M_{\theta,4} & M_{\zeta,4} + iM_{\eta,4} \\ 0 & 0 & M_{\zeta,4} - iM_{\eta,4} & M_{\theta,4} \\ -M_{\theta,4} & -M_{\zeta,4} - iM_{\eta,4} & 0 & 0 \\ -M_{\zeta,4} + iM_{\eta,4} & M_{\theta,4} & 0 & 0 \end{bmatrix}. \end{aligned}$$

Elements of these matrices can be turned by formula of shape [6]:

$$= \begin{pmatrix} \cos \frac{\theta}{2} & i \sin \frac{\theta}{2} \\ i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix} \begin{pmatrix} Z & X - iY \\ X + iY & -Z \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} & -i \sin \frac{\theta}{2} \\ -i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix}$$

$$= \begin{pmatrix} Z \cos \theta - Y \sin \theta & X - i(Y \cos \theta + Z \sin \theta) \\ X + i(Y \cos \theta + Z \sin \theta) & -Z \cos \theta + Y \sin \theta \end{pmatrix}.$$

Hence, if:

$$U_{2,3}(\alpha) \stackrel{def}{=} \begin{bmatrix} \cos \alpha & i \sin \alpha & 0 & 0 \\ i \sin \alpha & \cos \alpha & 0 & 0 \\ 0 & 0 & \cos \alpha & i \sin \alpha \\ 0 & 0 & i \sin \alpha & \cos \alpha \end{bmatrix}$$

and

$$\widehat{M}' \stackrel{def}{=} \begin{aligned} & -M'_{\zeta,0} \gamma_{\zeta}^{[0]} + M'_{\zeta,4} \zeta^{[4]} - \\ & -M'_{\eta,0} \gamma_{\eta}^{[0]} - M'_{\eta,4} \eta^{[4]} + \\ & + M'_{\theta,0} \gamma_{\theta}^{[0]} + M'_{\theta,4} \theta^{[4]} \end{aligned} \stackrel{def}{=} U_{2,3}^{\dagger}(\alpha) \widehat{M} U_{2,3}(\alpha)$$

then:

$$\begin{aligned} M'_{\zeta,0} &= M_{\zeta,0}, \\ M'_{\eta,0} &= M_{\eta,0} \cos 2\alpha + M_{\theta,0} \sin 2\alpha, \\ M'_{\theta,0} &= M_{\theta,0} \cos 2\alpha - M_{\eta,0} \sin 2\alpha, \\ M'_{\zeta,4} &= M_{\zeta,4}, \\ M'_{\eta,4} &= M_{\eta,4} \cos 2\alpha + M_{\theta,4} \sin 2\alpha, \\ M'_{\theta,4} &= M_{\theta,4} \cos 2\alpha - M_{\eta,4} \sin 2\alpha, \end{aligned}$$

Therefore, matrix $U_{2,3}(\alpha)$ makes an oscillation between green and blue quarks colors.

Let us consider equation (2) under transformation $U_{2,3}(\alpha)$ with α is an arbitrary real function of time-space variables ($\alpha = \alpha(t, x_1, x_2, x_3)$):

$$U_{2,3}^{\dagger}(\alpha) \left(\frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) U_{2,3}(\alpha) \varphi =$$

$$= U_{2,3}^{\dagger}(\alpha) \left(\begin{aligned} & \beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ & + \beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ & + \beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M} \end{aligned} \right) U_{2,3}(\alpha) \varphi.$$

That is:

$$\begin{pmatrix} \frac{1}{c} U_{2,3}^{\dagger}(\alpha) \partial_t (U_{2,3}(\alpha) \varphi) \\ + i\Theta_0 U_{2,3}^{\dagger}(\alpha) (U_{2,3}(\alpha) \varphi) \\ + i\Upsilon_0 U_{2,3}^{\dagger}(\alpha) \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{pmatrix} =$$

$$= U_{2,3}^\dagger(\alpha) \left(\begin{array}{l} \beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ + \beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ + \beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M} \end{array} \right) U_{2,3}(\alpha) \varphi.$$

Because

$$\begin{aligned} U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) &= 1_4, \\ U_{2,3}^\dagger(\alpha) \gamma^{[5]} U_{2,3}(\alpha) &= \gamma^{[5]} \end{aligned}$$

then

$$\begin{aligned} & \left(\frac{1}{c} \left(\left(U_{2,3}^\dagger(\alpha) \partial_t U_{2,3}(\alpha) \right) \varphi + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_t \varphi \right) + i\Theta_0 \varphi + i\Upsilon_0 \gamma^{[5]} \varphi \right) = \\ & = \left(\begin{array}{l} U_{2,3}^\dagger(\alpha) \beta^{[1]} \left(\begin{array}{l} (\partial_1 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_1 \varphi \\ + i\Theta_1 (U_{2,3}(\alpha) \varphi) + i\Upsilon_1 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{array} \right) + \\ + U_{2,3}^\dagger(\alpha) \beta^{[2]} \left(\begin{array}{l} (\partial_2 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_2 \varphi \\ + i\Theta_2 (U_{2,3}(\alpha) \varphi) + i\Upsilon_2 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{array} \right) + \\ + U_{2,3}^\dagger(\alpha) \beta^{[3]} \left(\begin{array}{l} (\partial_3 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_3 \varphi \\ + i\Theta_3 (U_{2,3}(\alpha) \varphi) + i\Upsilon_3 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \\ + iM_0 U_{2,3}^\dagger(\alpha) \gamma^{[0]} (U_{2,3}(\alpha) \varphi) \\ + iM_4 U_{2,3}^\dagger(\alpha) \beta^{[4]} (U_{2,3}(\alpha) \varphi) \\ + U_{2,3}^\dagger(\alpha) \widehat{M} (U_{2,3}(\alpha) \varphi) \end{array} \right) + \end{array} \right). \end{aligned}$$

Since

$$\begin{aligned} U_{2,3}^\dagger(\alpha) \gamma^{[0]} U_{2,3}(\alpha) &= \gamma^{[0]}, \\ U_{2,3}^\dagger(\alpha) \beta^{[4]} U_{2,3}(\alpha) &= \beta^{[4]} \end{aligned}$$

then

$$\begin{aligned} & \left(\frac{1}{c} \left(\left(U_{2,3}^\dagger(\alpha) \partial_t U_{2,3}(\alpha) \right) \varphi + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_t \varphi \right) + i\Theta_0 \varphi + i\Upsilon_0 \gamma^{[5]} \varphi \right) = \\ & = \left(\begin{array}{l} U_{2,3}^\dagger(\alpha) \beta^{[1]} \left(\begin{array}{l} (\partial_1 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_1 \varphi \\ + i\Theta_1 (U_{2,3}(\alpha) \varphi) + i\Upsilon_1 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{array} \right) + \\ + U_{2,3}^\dagger(\alpha) \beta^{[2]} \left(\begin{array}{l} (\partial_2 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_2 \varphi \\ + i\Theta_2 (U_{2,3}(\alpha) \varphi) + i\Upsilon_2 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{array} \right) + \\ + U_{2,3}^\dagger(\alpha) \beta^{[3]} \left(\begin{array}{l} (\partial_3 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_3 \varphi \\ + i\Theta_3 (U_{2,3}(\alpha) \varphi) + i\Upsilon_3 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \\ + iM_0 \gamma^{[0]} \varphi + iM_4 \beta^{[4]} \varphi + \widehat{M}' \varphi \end{array} \right) + \end{array} \right). \end{aligned}$$

Since

$$U_{2,3}^\dagger(\alpha) \beta^{[1]} = \beta^{[1]} U_{2,3}^\dagger(\alpha)$$

then

$$\begin{aligned} & \left(U_{2,3}^\dagger(\alpha) \frac{1}{c} \partial_t U_{2,3}(\alpha) + \frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi = \\ & = \left(\begin{aligned} & \beta^{[1]} U_{2,3}^\dagger(\alpha) \left(\begin{aligned} & (\partial_1 U_{2,3}(\alpha)) + U_{2,3}(\alpha) \partial_1 \\ & + i\Theta_1 U_{2,3}(\alpha) + i\Upsilon_1 \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + U_{2,3}^\dagger(\alpha) \beta^{[2]} \left(\begin{aligned} & (\partial_2 U_{2,3}(\alpha)) + U_{2,3}(\alpha) \partial_2 \\ & + i\Theta_2 U_{2,3}(\alpha) + i\Upsilon_2 \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + U_{2,3}^\dagger(\alpha) \beta^{[3]} \left(\begin{aligned} & (\partial_3 U_{2,3}(\alpha)) + U_{2,3}(\alpha) \partial_3 \\ & + i\Theta_3 U_{2,3}(\alpha) + i\Upsilon_3 \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M'} \end{aligned} \right) \varphi. \end{aligned}$$

Because

$$\begin{aligned} U_{2,3}^\dagger(\alpha) \beta^{[2]} &= (\beta^{[2]} \cos 2\alpha + \beta^{[3]} \sin 2\alpha) U_{2,3}^\dagger(\alpha), \\ U_{2,3}^\dagger(\alpha) \beta^{[3]} &= (\beta^{[3]} \cos 2\alpha - \beta^{[2]} \sin 2\alpha) U_{2,3}^\dagger(\alpha) \end{aligned}$$

then

$$\begin{aligned} & \left(U_{2,3}^\dagger(\alpha) \frac{1}{c} \partial_t U_{2,3}(\alpha) + \frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi = \\ & = \left(\begin{aligned} & \beta^{[1]} \left(\begin{aligned} & U_{2,3}^\dagger(\alpha) \partial_1 U_{2,3}(\alpha) + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_1 \\ & + i\Theta_1 U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) + i\Upsilon_1 U_{2,3}^\dagger(\alpha) \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + (\beta^{[2]} \cos 2\alpha + \beta^{[3]} \sin 2\alpha) \\ & \left(\begin{aligned} & U_{2,3}^\dagger(\alpha) \partial_2 U_{2,3}(\alpha) + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_2 \\ & + i\Theta_2 U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) + i\Upsilon_2 U_{2,3}^\dagger(\alpha) \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + (\beta^{[3]} \cos 2\alpha - \beta^{[2]} \sin 2\alpha) \\ & \left(\begin{aligned} & U_{2,3}^\dagger(\alpha) \partial_3 U_{2,3}(\alpha) + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_3 \\ & + i\Theta_3 U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) + i\Upsilon_3 U_{2,3}^\dagger(\alpha) \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M'} \end{aligned} \right) \varphi. \end{aligned}$$

Hence

$$\left(\frac{1}{c} \partial_t + U_{2,3}^\dagger(\alpha) \frac{1}{c} \partial_t U_{2,3}(\alpha) + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi =$$

$$= \left(\begin{array}{l} \beta^{[1]} \left(\partial_1 + U_{2,3}^\dagger(\alpha) \partial_1 U_{2,3}(\alpha) + i\Theta_1 + i\Upsilon_1 \gamma^{[5]} \right) \\ + \beta^{[2]} \left(\begin{array}{l} (\cos 2\alpha \cdot \partial_2 - \sin 2\alpha \cdot \partial_3) \\ + U_{2,3}^\dagger(\alpha) (\cos 2\alpha \cdot \partial_2 - \sin 2\alpha \cdot \partial_3) U_{2,3}(\alpha) \\ + i(\Theta_2 \cos 2\alpha - \Theta_3 \sin 2\alpha) \\ + i(\Upsilon_2 \gamma^{[5]} \cos 2\alpha - \Upsilon_3 \gamma^{[5]} \sin 2\alpha) \end{array} \right) \\ + \beta^{[3]} \left(\begin{array}{l} (\cos 2\alpha \cdot \partial_3 + \sin 2\alpha \cdot \partial_2) \\ + U_{2,3}^\dagger(\alpha) (\cos 2\alpha \cdot \partial_3 + \sin 2\alpha \cdot \partial_2) U_{2,3}(\alpha) \\ + i(\Theta_2 \sin 2\alpha + \Theta_3 \cos 2\alpha) \\ + i(\Upsilon_3 \gamma^{[5]} \cos 2\alpha + \Upsilon_2 \gamma^{[5]} \sin 2\alpha) \end{array} \right) + \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}' \end{array} \right) \varphi. \quad (3)$$

Let x'_2 and x'_3 are elements of other coordinates system such that:

$$\begin{aligned} \frac{\partial x_2}{\partial x'_2} &= \cos 2\alpha, \\ \frac{\partial x_3}{\partial x'_2} &= -\sin 2\alpha, \\ \frac{\partial x_2}{\partial x'_3} &= \sin 2\alpha, \\ \frac{\partial x_3}{\partial x'_3} &= \cos 2\alpha, \\ \frac{\partial x_0}{\partial x'_2} &= \frac{\partial x_1}{\partial x'_2} = \frac{\partial x_0}{\partial x'_3} = \frac{\partial x_1}{\partial x'_3} = 0. \end{aligned}$$

Hence:

$$\begin{aligned} \partial_2' \stackrel{def}{=} \frac{\partial}{\partial x'_2} &= \frac{\partial}{\partial x_0} \frac{\partial x_0}{\partial x'_2} + \frac{\partial}{\partial x_1} \frac{\partial x_1}{\partial x'_2} + \frac{\partial}{\partial x_2} \frac{\partial x_2}{\partial x'_2} + \frac{\partial}{\partial x_3} \frac{\partial x_3}{\partial x'_2} \\ &= \cos 2\alpha \cdot \frac{\partial}{\partial x_2} - \sin 2\alpha \cdot \frac{\partial}{\partial x_3} \\ &= \cos 2\alpha \cdot \partial_2 - \sin 2\alpha \cdot \partial_3, \\ \\ \partial_3' \stackrel{def}{=} \frac{\partial}{\partial x'_3} &= \frac{\partial}{\partial x_0} \frac{\partial x_0}{\partial x'_3} + \frac{\partial}{\partial x_1} \frac{\partial x_1}{\partial x'_3} + \frac{\partial}{\partial x_2} \frac{\partial x_2}{\partial x'_3} + \frac{\partial}{\partial x_3} \frac{\partial x_3}{\partial x'_3} \\ &= \cos 2\alpha \cdot \frac{\partial}{\partial x_3} + \sin 2\alpha \cdot \frac{\partial}{\partial x_2} \\ &= \cos 2\alpha \cdot \partial_3 + \sin 2\alpha \cdot \partial_2. \end{aligned}$$

Therefore from (3):

$$\begin{aligned}
& \left(\frac{1}{c} \partial_t + U_{2,3}^\dagger(\alpha) \frac{1}{c} \partial_t U_{2,3}(\alpha) + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi = \\
& = \left(\begin{aligned} & \beta^{[1]} \left(\partial_1 + U_{2,3}^\dagger(\alpha) \partial_1 U_{2,3}(\alpha) + i\Theta_1 + i\Upsilon_1 \gamma^{[5]} \right) \\ & + \beta^{[2]} \left(\partial'_2 + U_{2,3}^\dagger(\alpha) \partial'_2 U_{2,3}(\alpha) + i\Theta'_2 + i\Upsilon'_2 \gamma^{[5]} \right) \\ & + \beta^{[3]} \left(\partial'_3 + U_{2,3}^\dagger(\alpha) \partial'_3 U_{2,3}(\alpha) + i\Theta'_3 + i\Upsilon'_3 \gamma^{[5]} \right) + \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}' \end{aligned} \right) \varphi.
\end{aligned}$$

with

$$\begin{aligned}
\Theta'_2 &\stackrel{def}{=} \Theta_2 \cos 2\alpha - \Theta_3 \sin 2\alpha, \\
\Theta'_3 &\stackrel{def}{=} \Theta_2 \sin 2\alpha + \Theta_3 \cos 2\alpha. \\
\Upsilon'_2 &\stackrel{def}{=} \Upsilon_2 \cos 2\alpha - \Upsilon_3 \sin 2\alpha, \\
\Upsilon'_3 &\stackrel{def}{=} \Upsilon_2 \sin 2\alpha + \Upsilon_3 \cos 2\alpha.
\end{aligned}$$

Therefore, the oscillation between blue and green quarks colors curves the space of events in the x_2, x_3 directions.

Similarly that: matrix

$$U_{1,3}(\vartheta) \stackrel{def}{=} \begin{bmatrix} \cos \vartheta & \sin \vartheta & 0 & 0 \\ -\sin \vartheta & \cos \vartheta & 0 & 0 \\ 0 & 0 & \cos \vartheta & \sin \vartheta \\ 0 & 0 & -\sin \vartheta & \cos \vartheta \end{bmatrix}$$

with an arbitrary real function $\vartheta(t, x_1, x_2, x_3)$ describes the oscillation between blue and red quarks colors which curves the space of events in the x_1, x_3 directions. And matrix

$$U_{1,2}(\varsigma) \stackrel{def}{=} \begin{bmatrix} \cos \varsigma - i \sin \varsigma & 0 & 0 & 0 \\ 0 & \cos \varsigma + i \sin \varsigma & 0 & 0 \\ 0 & 0 & \cos \varsigma - i \sin \varsigma & 0 \\ 0 & 0 & 0 & \cos \varsigma + i \sin \varsigma \end{bmatrix}$$

with an arbitrary real function $\varsigma(t, x_1, x_2, x_3)$ describes the oscillation between green and red quarks colors which curves the space of events in the x_1, x_2 directions.

Now, let

$$U_{0,1}(v) \stackrel{def}{=} \begin{bmatrix} \cosh v & -\sinh v & 0 & 0 \\ -\sinh v & \cosh v & 0 & 0 \\ 0 & 0 & \cosh v & \sinh v \\ 0 & 0 & \sinh v & \cosh v \end{bmatrix}.$$

and

$$\widehat{M}'' \stackrel{def}{=} \begin{aligned} & -M''_{\zeta,0}\gamma_{\zeta}^{[0]} + M''_{\zeta,4}\zeta^{[4]} - \\ & -M''_{\eta,0}\gamma_{\eta}^{[0]} - M''_{\eta,4}\eta^{[4]} + \\ & + M''_{\theta,0}\gamma_{\theta}^{[0]} + M''_{\theta,4}\theta^{[4]} \end{aligned} \stackrel{def}{=} U_{0,1}^{\dagger}(v) \widehat{M} U_{0,1}(v)$$

then:

$$\begin{aligned} M''_{\zeta,0} &= M_{\zeta,0}, \\ M''_{\eta,0} &= M_{\eta,0} \cosh 2v - M_{\theta,4} \sinh 2v, \\ M''_{\theta,0} &= M_{\theta,0} \cosh 2v + M_{\eta,4} \sinh 2v, \\ M''_{\zeta,4} &= M_{\zeta,4} \\ M''_{\eta,4} &= M_{\eta,4} \cosh 2v + M_{\theta,0} \sinh 2v, \\ M''_{\theta,4} &= M_{\theta,4} \cosh 2v - M_{\eta,0} \sinh 2v, \end{aligned}$$

Therefore, matrix $U_{0,1}(v)$ makes an oscillation between green and blue quarks colors with an oscillation between up and down quarks.

Let us consider equation (2) under transformation $U_{0,1}(v)$ with v is an arbitrary real function of time-space variables ($v = v(t, x_1, x_2, x_3)$):

$$\begin{aligned} & U_{0,1}^{\dagger}(v) \left(\frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) U_{0,1}(v) \varphi = \\ & = U_{0,1}^{\dagger}(v) \left(\begin{aligned} & \beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ & + \beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ & + \beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M} \end{aligned} \right) U_{0,1}(v) \varphi. \end{aligned}$$

Hence,

$$\begin{aligned} & U_{0,1}^{\dagger}(v) \left(\begin{aligned} & \frac{1}{c} \partial_t (U_{0,1}(v) \varphi) \\ & + i\Theta_0 (U_{0,1}(v) \varphi) + i\Upsilon_0 \gamma^{[5]} (U_{0,1}(v) \varphi) \end{aligned} \right) = \\ & = U_{0,1}^{\dagger}(v) \left(\begin{aligned} & \beta^{[1]} \left(\begin{aligned} & \partial_1 (U_{0,1}(v) \varphi) \\ & + i\Theta_1 (U_{0,1}(v) \varphi) + i\Upsilon_1 \gamma^{[5]} (U_{0,1}(v) \varphi) \end{aligned} \right) + \\ & + \beta^{[2]} \left(\begin{aligned} & \partial_2 (U_{0,1}(v) \varphi) \\ & + i\Theta_2 (U_{0,1}(v) \varphi) + i\Upsilon_2 \gamma^{[5]} (U_{0,1}(v) \varphi) \end{aligned} \right) + \\ & + \beta^{[3]} \left(\begin{aligned} & \partial_3 (U_{0,1}(v) \varphi) \\ & + i\Theta_3 (U_{0,1}(v) \varphi) + i\Upsilon_3 \gamma^{[5]} (U_{0,1}(v) \varphi) \end{aligned} \right) + \\ & + iM_0 \gamma^{[0]} (U_{0,1}(v) \varphi) \\ & + iM_4 \beta^{[4]} (U_{0,1}(v) \varphi) + \widehat{M} (U_{0,1}(v) \varphi) \end{aligned} \right). \\ & U_{0,1}^{\dagger}(v) \left(\begin{aligned} & \frac{1}{c} (\partial_t U_{0,1}(v)) \varphi + U_{0,1}(v) \frac{1}{c} \partial_t \varphi \\ & + i\Theta_0 U_{0,1}(v) \varphi + i\Upsilon_0 \gamma^{[5]} U_{0,1}(v) \varphi \end{aligned} \right) = \end{aligned}$$

$$= U_{0,1}^\dagger(v) \left(\begin{array}{l} \beta^{[1]} \left(\begin{array}{l} (\partial_1 U_{0,1}(v)) \varphi + U_{0,1}(v) \partial_1 \varphi \\ + i\Theta_1 U_{0,1}(v) \varphi + i\Upsilon_1 \gamma^{[5]} U_{0,1}(v) \varphi \end{array} \right) + \\ + \beta^{[2]} \left(\begin{array}{l} (\partial_2 U_{0,1}(v)) \varphi + U_{0,1}(v) \partial_2 \varphi \\ + i\Theta_2 U_{0,1}(v) \varphi + i\Upsilon_2 \gamma^{[5]} U_{0,1}(v) \varphi \end{array} \right) + \\ + \beta^{[3]} \left(\begin{array}{l} (\partial_3 U_{0,1}(v)) \varphi + U_{0,1}(v) \partial_3 \varphi \\ + i\Theta_3 U_{0,1}(v) \varphi + i\Upsilon_3 \gamma^{[5]} U_{0,1}(v) \varphi \end{array} \right) + \\ + iM_0 \gamma^{[0]} U_{0,1}(v) \varphi \\ + iM_4 \beta^{[4]} U_{0,1}(v) \varphi + \widehat{M} U_{0,1}(v) \varphi \end{array} \right).$$

Therefore:

$$\left(\begin{array}{l} U_{0,1}^\dagger(v) \frac{1}{c} (\partial_t U_{0,1}(v)) + U_{0,1}^\dagger(v) U_{0,1}(v) \frac{1}{c} \partial_t \\ + i\Theta_0 U_{0,1}^\dagger(v) U_{0,1}(v) + i\Upsilon_0 U_{0,1}^\dagger(v) \gamma^{[5]} U_{0,1}(v) \end{array} \right) \varphi =$$

$$= \left(\begin{array}{l} U_{0,1}^\dagger(v) \beta^{[1]} \left(\begin{array}{l} (\partial_1 U_{0,1}(v)) + U_{0,1}(v) \partial_1 \\ + i\Theta_1 U_{0,1}(v) + i\Upsilon_1 \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + U_{0,1}^\dagger(v) \beta^{[2]} \left(\begin{array}{l} (\partial_2 U_{0,1}(v)) + U_{0,1}(v) \partial_2 \\ + i\Theta_2 U_{0,1}(v) + i\Upsilon_2 \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + U_{0,1}^\dagger(v) \beta^{[3]} \left(\begin{array}{l} (\partial_3 U_{0,1}(v)) + U_{0,1}(v) \partial_3 \\ + i\Theta_3 U_{0,1}(v) + i\Upsilon_3 \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + iM_0 U_{0,1}^\dagger(v) \gamma^{[0]} U_{0,1}(v) \\ + iM_4 U_{0,1}^\dagger(v) \beta^{[4]} U_{0,1}(v) + U_{0,1}^\dagger(v) \widehat{M} U_{0,1}(v) \end{array} \right) \varphi.$$

Since (about a sum of matrix and number in [8]):

$$\begin{aligned} U_{0,1}^\dagger(v) U_{0,1}(v) &= (\cosh 2v - \beta^{[1]} \sinh 2v), \\ U_{0,1}^\dagger(v) &= (\cosh 2v + \beta^{[1]} \sinh 2v) U_{0,1}^{-1}(v), \\ U_{0,1}^\dagger(v) \beta^{[1]} &= (\beta^{[1]} \cosh 2v - \sinh 2v) U_{0,1}^{-1}(v), \\ U_{0,1}^\dagger(v) \beta^{[2]} &= \beta^{[2]} U_{0,1}^{-1}(v), \\ U_{0,1}^\dagger(v) \beta^{[3]} &= \beta^{[3]} U_{0,1}^{-1}(v), \\ U_{0,1}^\dagger(v) \gamma^{[0]} U_{0,1}(v) &= \gamma^{[0]}, \\ U_{0,1}^\dagger(v) \beta^{[4]} U_{0,1}(v) &= \beta^{[4]} \end{aligned}$$

then

$$\left(\begin{array}{l} (\cosh 2v + \beta^{[1]} \sinh 2v) U_{0,1}^{-1}(v) \frac{1}{c} (\partial_t U_{0,1}(v)) \\ + (\cosh 2v - \beta^{[1]} \sinh 2v) \frac{1}{c} \partial_t \\ + i\Theta_0 (\cosh 2v - \beta^{[1]} \sinh 2v) \\ + i\Upsilon_0 U_{0,1}^\dagger(v) \gamma^{[5]} U_{0,1}(v) \end{array} \right) \varphi =$$

$$= \left(\begin{array}{c} (\beta^{[1]} \cosh 2v - \sinh 2v) \cdot \\ \cdot U_{0,1}^{-1}(v) \left(\begin{array}{c} (\partial_1 U_{0,1}(v)) + U_{0,1}(v) \partial_1 \\ + i\Theta_1 U_{0,1}(v) + i\Upsilon_1 \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + \beta^{[2]} U_{0,1}^{-1}(v) \left(\begin{array}{c} (\partial_2 U_{0,1}(v)) + U_{0,1}(v) \partial_2 \\ + i\Theta_2 U_{0,1}(v) + i\Upsilon_2 \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + \beta^{[3]} U_{0,1}^{-1}(v) \left(\begin{array}{c} (\partial_3 U_{0,1}(v)) + U_{0,1}(v) \partial_3 \\ + i\Theta_3 U_{0,1}(v) + i\Upsilon_3 \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}'' \end{array} \right) \varphi.$$

Hence:

$$\left(\begin{array}{c} (\cosh 2v + \beta^{[1]} \sinh 2v) U_{0,1}^{-1}(v) \frac{1}{c} (\partial_t U_{0,1}(v)) \\ + (\cosh 2v - \beta^{[1]} \sinh 2v) \frac{1}{c} \partial_t \\ + i\Theta_0 (\cosh 2v - \beta^{[1]} \sinh 2v) \\ + i\Upsilon_0 U_{0,1}^\dagger(v) \gamma^{[5]} U_{0,1}(v) \end{array} \right) \varphi =$$

$$= \left(\begin{array}{c} (\beta^{[1]} \cosh 2v - \sinh 2v) \cdot \\ \cdot \left(\begin{array}{c} U_{0,1}^{-1}(v) (\partial_1 U_{0,1}(v)) + U_{0,1}^{-1}(v) U_{0,1}(v) \partial_1 \\ + i\Theta_1 U_{0,1}^{-1}(v) U_{0,1}(v) + i\Upsilon_1 U_{0,1}^{-1}(v) \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + \beta^{[2]} \left(\begin{array}{c} U_{0,1}^{-1}(v) (\partial_2 U_{0,1}(v)) + U_{0,1}^{-1}(v) U_{0,1}(v) \partial_2 \\ + i\Theta_2 U_{0,1}^{-1}(v) U_{0,1}(v) + i\Upsilon_2 U_{0,1}^{-1}(v) \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + \beta^{[3]} \left(\begin{array}{c} U_{0,1}^{-1}(v) (\partial_3 U_{0,1}(v)) + U_{0,1}^{-1}(v) U_{0,1}(v) \partial_3 \\ + i\Theta_3 U_{0,1}^{-1}(v) U_{0,1}(v) + i\Upsilon_3 U_{0,1}^{-1}(v) \gamma^{[5]} U_{0,1}(v) \end{array} \right) + \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}'' \end{array} \right) \varphi.$$

Because

$$\begin{aligned} U_{0,1}^{-1}(v) U_{0,1}(v) &= 1_4, \\ U_{0,1}^{-1}(v) \gamma^{[5]} U_{0,1}(v) &= \gamma^{[5]}, \\ U_{0,1}^\dagger(v) \gamma^{[5]} U_{0,1}(v) &= \gamma^{[5]} (\cosh 2v - \beta^{[1]} \sinh 2v) \end{aligned}$$

then

$$\left(\begin{array}{c} (\cosh 2v + \beta^{[1]} \sinh 2v) U_{0,1}^{-1}(v) \frac{1}{c} \partial_t U_{0,1}(v) \\ + (\cosh 2v - \beta^{[1]} \sinh 2v) \frac{1}{c} \partial_t \\ + i\Theta_0 (\cosh 2v - \beta^{[1]} \sinh 2v) \\ + i\Upsilon_0 \gamma^{[5]} (\cosh 2v - \beta^{[1]} \sinh 2v) \end{array} \right) \varphi =$$

$$= \left(\begin{array}{c} (\beta^{[1]} \cosh 2v - \sinh 2v) \cdot \\ \cdot (\partial_1 + U_{0,1}^{-1}(v) (\partial_1 U_{0,1}(v)) + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ + \beta^{[2]} (\partial_2 + U_{0,1}^{-1}(v) (\partial_2 U_{0,1}(v)) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ + \beta^{[3]} (\partial_3 + U_{0,1}^{-1}(v) (\partial_3 U_{0,1}(v)) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}'' \end{array} \right) \varphi.$$

Hence:

$$\left(\begin{array}{l} \cosh 2v \cdot U_{0,1}^{-1}(v) \frac{1}{c} \partial_t U_{0,1}(v) + \cosh 2v \cdot \frac{1}{c} \partial_t + \sinh 2v \cdot \partial_1 \\ + i\Theta_0 \cosh 2v + i\Upsilon_0 \gamma^{[5]} \cosh 2v \\ + \sinh 2v \cdot U_{0,1}^{-1}(v) (\partial_1 U_{0,1}(v)) \\ + i\Theta_1 \sinh 2v \\ + i \sinh 2v \cdot \Upsilon_1 \gamma^{[5]} \\ + \beta^{[1]} \sinh 2v \cdot U_{0,1}^{-1}(v) \frac{1}{c} \partial_t U_{0,1}(v) - \beta^{[1]} \sinh 2v \cdot \frac{1}{c} \partial_t \\ - i\beta^{[1]} \Theta_0 \sinh 2v - i\beta^{[1]} \Upsilon_0 \gamma^{[5]} \sinh 2v - \beta^{[1]} \cosh 2v \cdot \partial_1 \\ - \beta^{[1]} \cosh 2v \cdot U_{0,1}^{-1}(v) (\partial_1 U_{0,1}(v)) \\ - i\beta^{[1]} \Theta_1 \cosh 2v \\ - i\beta^{[1]} \cosh 2v \cdot \Upsilon_1 \gamma^{[5]} \\ - \beta^{[2]} (\partial_2 + U_{0,1}^{-1}(v) (\partial_2 U_{0,1}(v)) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) \\ - \beta^{[3]} (\partial_3 + U_{0,1}^{-1}(v) (\partial_3 U_{0,1}(v)) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) \\ - iM_0 \gamma^{[0]} - iM_4 \beta^{[4]} - \widehat{M}'' \end{array} \right) \varphi = 0.$$

Therefore,

$$\left(\begin{array}{l} U_{0,1}^{-1}(v) (\cosh 2v \cdot \frac{1}{c} \partial_t + \sinh 2v \cdot \partial_1) U_{0,1}(v) \\ + (\cosh 2v \cdot \frac{1}{c} \partial_t + \sinh 2v \cdot \partial_1) \\ + i (\Theta_0 \cosh 2v + \Theta_1 \sinh 2v) \\ + i (\Upsilon_0 \cosh 2v + \sinh 2v \cdot \Upsilon_1) \gamma^{[5]} \\ - \beta^{[1]} \left(\begin{array}{l} U_{0,1}^{-1}(v) (\cosh 2v \cdot \partial_1 + \sinh 2v \cdot \frac{1}{c} \partial_t) U_{0,1}(v) \\ + (\cosh 2v \cdot \partial_1 + \sinh 2v \cdot \frac{1}{c} \partial_t) \\ + i (\Theta_1 \cosh 2v + \Theta_0 \sinh 2v) \\ + i (\Upsilon_1 \cosh 2v + \Upsilon_0 \sinh 2v) \gamma^{[5]} \end{array} \right) \\ - \beta^{[2]} (\partial_2 + U_{0,1}^{-1}(v) (\partial_2 U_{0,1}(v)) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) \\ - \beta^{[3]} (\partial_3 + U_{0,1}^{-1}(v) (\partial_3 U_{0,1}(v)) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) \\ - iM_0 \gamma^{[0]} - iM_4 \beta^{[4]} - \widehat{M}'' \end{array} \right) \varphi = 0 \quad (4)$$

Let t' and x'_1 are elements of other coordinates system such that:

$$\begin{aligned} \frac{\partial x_1}{\partial x'_1} &= \cosh 2v, \\ \frac{\partial t}{\partial x'_1} &= \frac{1}{c} \sinh 2v, \\ \frac{\partial x_1}{\partial t'} &= c \sinh 2v. \\ \frac{\partial t}{\partial t'} &= \cosh 2v, \\ \frac{\partial x_2}{\partial t'} &= \frac{\partial x_3}{\partial t'} = \frac{\partial x_2}{\partial x'_1} = \frac{\partial x_3}{\partial x'_1} = 0. \end{aligned}$$

Hence:

$$\begin{aligned}
\partial_t' \stackrel{def}{=} \frac{\partial}{\partial t'} &= \frac{\partial}{\partial t} \frac{\partial t}{\partial t'} + \frac{\partial}{\partial x_1} \frac{\partial x_1}{\partial t'} + \frac{\partial}{\partial x_2} \frac{\partial x_2}{\partial t'} + \frac{\partial}{\partial x_3} \frac{\partial x_3}{\partial t'} \\
&= \cosh 2v \cdot \frac{\partial}{\partial t} + c \sinh 2v \cdot \frac{\partial}{\partial x_1} \\
&= \cosh 2v \cdot \partial_t + c \sinh 2v \cdot \partial_1,
\end{aligned}$$

That is:

$$\frac{1}{c} \partial_t' = \frac{1}{c} \cosh 2v \cdot \partial_t + \sinh 2v \cdot \partial_1.$$

And

$$\begin{aligned}
\partial_1' \stackrel{def}{=} \frac{\partial}{\partial x_1'} &= \frac{\partial}{\partial t} \frac{\partial t}{\partial x_1'} + \frac{\partial}{\partial x_1} \frac{\partial x_1}{\partial x_1'} + \frac{\partial}{\partial x_2} \frac{\partial x_2}{\partial x_1'} + \frac{\partial}{\partial x_3} \frac{\partial x_3}{\partial x_1'} \\
&= \cosh 2v \cdot \frac{\partial}{\partial x_1} + \sinh 2v \cdot \frac{1}{c} \frac{\partial}{\partial t} \\
&= \cosh 2v \cdot \partial_1 + \sinh 2v \cdot \frac{1}{c} \partial_t.
\end{aligned}$$

Therefore from (4):

$$\begin{pmatrix}
+\frac{1}{c} \partial_t' + U_{0,1}^{-1}(v) \frac{1}{c} \partial_t' U_{0,1}(v) + i\Theta_0'' + i\Upsilon_0'' \gamma^{[5]} \\
-\beta^{[1]} (+\partial_1' + U_{0,1}^{-1}(v) \partial_1' U_{0,1}(v) + i\Theta_1'' + i\Upsilon_1'' \gamma^{[5]}) \\
-\beta^{[2]} (\partial_2 + U_{0,1}^{-1}(v) (\partial_2 U_{0,1}(v)) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) \\
-\beta^{[3]} (\partial_3 + U_{0,1}^{-1}(v) (\partial_3 U_{0,1}(v)) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) \\
-iM_0 \gamma^{[0]} - iM_4 \beta^{[4]} - \widehat{M}''
\end{pmatrix} \varphi = 0$$

with

$$\begin{aligned}
\Theta_0'' &\stackrel{def}{=} \Theta_0 \cosh 2v + \Theta_1 \sinh 2v, \\
\Theta_1'' &\stackrel{def}{=} \Theta_1 \cosh 2v + \Theta_0 \sinh 2v, \\
\Upsilon_0'' &\stackrel{def}{=} \Upsilon_0 \cosh 2v + \sinh 2v \cdot \Upsilon_1, \\
\Upsilon_1'' &\stackrel{def}{=} \Upsilon_1 \cosh 2v + \Upsilon_0 \sinh 2v.
\end{aligned}$$

Therefore, the oscillation between blue and green quarks colors with the oscillation between up and down quarks curves the space of events in the t , x_1 directions.

Similarly that: matrix

$$U_{0,2}(\phi) = \begin{bmatrix} \cosh \phi & i \sinh \phi & 0 & 0 \\ -i \sinh \phi & \cosh \phi & 0 & 0 \\ 0 & 0 & \cosh \phi & -i \sinh \phi \\ 0 & 0 & i \sinh \phi & \cosh \phi \end{bmatrix}$$

with an arbitrary real function $\phi(t, x_1, x_2, x_3)$ describes the oscillation between blue and red quarks colors with the oscillation between up and down quarks which curves the space of events in the t, x_2 directions. And matrix

$$U_{0,3}(\iota) = \begin{bmatrix} \cosh \iota + \sinh \iota & 0 & 0 & 0 \\ 0 & \cosh \iota - \sinh \iota & 0 & 0 \\ 0 & 0 & \cosh \iota - \sinh \iota & 0 \\ 0 & 0 & 0 & \cosh \iota + \sinh \iota \end{bmatrix}$$

with an arbitrary real function $\iota(t, x_1, x_2, x_3)$ describes the oscillation between green and red quarks colors with the oscillation between up and down quarks which curves the space of events in the t, x_3 directions.

Now, because

$$\iota(t, x_1, x_2, x_3), \phi(t, x_1, x_2, x_3), v = v(t, x_1, x_2, x_3), \varsigma(t, x_1, x_2, x_3), \\ \vartheta(t, x_1, x_2, x_3), \alpha(t, x_1, x_2, x_3)$$

are arbitrary real functions then for coordinates X_ν of the curved time-space the formula 2, page 155 from [7] is fulfils:

$$dX_\nu = \sum_{\sigma=0}^3 a_{\nu\sigma} dx_\sigma$$

with real functions $a_{\nu\sigma}(t, x_1, x_2, x_3)$.

Hence, the square of the linear element ds^2 obeys to formula 3, page 155:

$$ds^2 = \sum_{\sigma=0}^3 \sum_{\tau=0}^3 g_{\sigma\tau} dx_\sigma dx_\tau$$

with real functions $g_{\sigma\tau}(t, x_1, x_2, x_3)$ such that $g_{\sigma\tau} = g_{\tau\sigma}$. And then the Einstein deductions from [7] can be reiterated:

The equation of a geodesic line between points P and P' is the following (formula 20, page 167):

$$\delta \int_P^{P'} ds = 0.$$

Hence with a parameter λ :

$$\delta \int_{\lambda_1}^{\lambda_2} \sqrt{\sum_{\sigma=0}^3 \sum_{\tau=0}^3 g_{\sigma\tau} \frac{dx_\sigma}{d\lambda} \frac{dx_\tau}{d\lambda}} d\lambda = 0.$$

If denote:

$$F(x_\sigma, x_\tau, \frac{dx_\sigma}{d\lambda}, \frac{dx_\tau}{d\lambda}) \stackrel{def}{=} \sqrt{\sum_{\sigma=0}^3 \sum_{\tau=0}^3 g_{\sigma\tau} \frac{dx_\sigma}{d\lambda} \frac{dx_\tau}{d\lambda}}$$

then in accordance with the Euler-Lagrange variational principle the geodesic line equation is the following:

$$\frac{\partial F}{\partial x_\sigma} - \frac{d}{d\lambda} \frac{\partial F}{\partial \left(\frac{dx_\sigma}{d\lambda}\right)} = 0.$$

Because

$$\begin{aligned} \frac{\partial F}{\partial x_\sigma} &= \frac{\partial}{\partial x_\sigma} \left(\sqrt{\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda}} \right) \\ &= \frac{1}{2} \frac{1}{\sqrt{\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda}}} \sum_{\nu=0}^3 \sum_{\mu=0}^3 \frac{\partial g_{\nu\mu}}{\partial x_\sigma} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda} \end{aligned}$$

and

$$\begin{aligned} \frac{d}{d\lambda} \frac{\partial F}{\partial \left(\frac{dx_\sigma}{d\lambda}\right)} &= \frac{d}{d\lambda} \frac{\partial}{\partial \left(\frac{dx_\sigma}{d\lambda}\right)} \left(\sqrt{\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda}} \right) \\ &= \frac{d}{d\lambda} \frac{1}{\sqrt{\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda}}} \sum_{\mu=0}^3 g_{\sigma\mu} \frac{dx_\mu}{d\lambda}. \end{aligned}$$

then the equation of geodesic line is the following:

$$\begin{aligned} &\frac{1}{2} \frac{1}{\sqrt{\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda}}} \sum_{\nu=0}^3 \sum_{\mu=0}^3 \frac{\partial g_{\nu\mu}}{\partial x_\sigma} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda} \\ &- \frac{d}{d\lambda} \frac{1}{\sqrt{\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda}}} \sum_{\mu=0}^3 g_{\sigma\mu} \frac{dx_\mu}{d\lambda} = 0. \end{aligned}$$

if $\lambda = s$ then

$$\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda} = \frac{\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} dx_\nu dx_\mu}{d\lambda^2} = \frac{ds^2}{ds^2} = 1,$$

Hence

$$\frac{d}{d\lambda} \frac{1}{\sqrt{\sum_{\nu=0}^3 \sum_{\mu=0}^3 g_{\nu\mu} \frac{dx_\nu}{d\lambda} \frac{dx_\mu}{d\lambda}}} \sum_{\mu=0}^3 g_{\sigma\mu} \frac{dx_\mu}{d\lambda} = \frac{d}{d\lambda} \sum_{\mu=0}^3 g_{\sigma\mu} \frac{dx_\mu}{d\lambda}$$

$$\begin{aligned}
&= \sum_{\mu=0}^3 \left(\left(\frac{d}{d\lambda} g_{\sigma\mu} \right) \frac{dx_\mu}{d\lambda} + g_{\sigma\mu} \frac{d}{d\lambda} \frac{dx_\mu}{d\lambda} \right) \\
&= \sum_{\mu=0}^3 \left(\sum_{\tau=0}^3 \frac{\partial g_{\sigma\mu}}{\partial x_\tau} \frac{\partial x_\tau}{d\lambda} \frac{dx_\mu}{d\lambda} + g_{\sigma\mu} \frac{d}{d\lambda} \frac{dx_\mu}{d\lambda} \right) \\
&= \sum_{\mu=0}^3 \left(\sum_{\tau=0}^3 \frac{\partial g_{\sigma\mu}}{\partial x_\tau} \frac{\partial x_\tau}{ds} \frac{dx_\mu}{ds} + g_{\sigma\mu} \frac{d}{ds} \frac{dx_\mu}{ds} \right).
\end{aligned}$$

Therefore, the geodesic line equation is the following:

$$\sum_{\mu=0}^3 \left(g_{\sigma\mu} \frac{d^2 x_\mu}{ds^2} + \sum_{\tau=1}^3 \frac{\partial g_{\sigma\mu}}{\partial x_\tau} \frac{\partial x_\tau}{ds} \frac{dx_\mu}{ds} \right) - \frac{1}{2} \sum_{\mu=0}^3 \sum_{\nu=0}^3 \frac{\partial g_{\nu\mu}}{\partial x_\sigma} \frac{dx_\nu}{ds} \frac{dx_\mu}{ds} = 0.$$

=====

References

- [1] G. Quznetsov, *preprint* arXiv:0708.2322v1
- [2] G. Quznetsov, *Probabilistic Treatment of Gauge Theories*, in series *Contemporary Fundamental Physics*, ed. V. Dvoeglazov, Nova Sci., NY (2007) pp.38-39
- [3] Item, p.12
- [4] Item, pp.95-117
- [5] Item, p.p.68-72
- [6] For instance J. M. Ziman, *Elements of Advanced Quantum Theory*, Cambridge University Press (1969) formula (6.59)
- [7] Albert Einstein, *The Foundation of the General Theory of Relativity*, Princeton University Press (1997) <http://www.alberteinstein.info/gallery/pdf/>
- [8] G. Quznetsov, *Probabilistic Treatment of Gauge Theories*, in series *Contemporary Fundamental Physics*, ed. V. Dvoeglazov, Nova Sci., NY (2007) p.13